

8-28-2017

## Problem Set #1

David Peak  
*Utah State University*, [david.peak@usu.edu](mailto:david.peak@usu.edu)

Follow this and additional works at: [https://digitalcommons.usu.edu/intro\\_modernphysics\\_problems](https://digitalcommons.usu.edu/intro_modernphysics_problems)



Part of the [Physics Commons](#)

---

### Recommended Citation

Peak, David, "Problem Set #1" (2017). *Problems*. Paper 1.  
[https://digitalcommons.usu.edu/intro\\_modernphysics\\_problems/1](https://digitalcommons.usu.edu/intro_modernphysics_problems/1)

This Course is brought to you for free and open access by the Introductory Modern Physics at DigitalCommons@USU. It has been accepted for inclusion in Problems by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



Problems 1-2 refer to: The electric field in a laser beam is given by

$$\mathcal{E}(x,t) = (1000\text{V/m})\sin\left[\left(\pi\times 10^7\text{rad/m}\right)x + \left(3\pi\times 10^{15}\text{rad/s}\right)t\right].$$

1. Answer these questions by referring to the general form for a plane wave found on Fn1, p3.  
(a) What is the wavelength of the beam? Is the laser beam visible light, IR, or UV? Why? (b) What is the (ordinary) frequency of the beam? (c) Using the results of (a) and (b) verify that the expression given above describes a light wave. (d) How long does it take the beam to travel 1 nm? (e) In which direction is the beam propagating? Explain.
2. Given that the magnitude of the electric and magnetic fields in the laser beam are related by  $\mathcal{E} = cB$ , find the maximum value of the energy density  $u$ .
3. A double slit interference pattern appears on a whiteboard 1 m from the plate with holes. The wavelength of the light is 550 nm. The distance from the central maximum to the first adjacent maximum is 5 cm. What is the slit separation in the plate? In principle, how many maxima might be observed? Explain all steps.